

## **DALACHOY TARKIBIDAGI FLAVONOIDLAR MIQDORINI ANIQLASH**

**Ibrohimjon ASQAROV,**  
**Andijon davlat universiteti**  
**Kimyo kafedrası professori,**  
**O‘zbekistonda xizmat ko‘rsatgan ixtirochi,**  
**O‘zbekiston TABOBAT Akademiyasi raisi**  
**Go‘yoxon MO‘MINOVA,**  
**Andijon davlat tibbiyot instituti**  
**Biologik kimyo kafedrası dotsenti,**  
**biologiya fanlari bo'yicha falsafa doktori (PhD)**  
**E-mail:([guvoxonmuminova@gmail.com](mailto:guvoxonmuminova@gmail.com)).**

**Annotatsiya:** Dalachoy o‘simligi tarkibi o‘rganilganda flavonoidlarga boy ekani aniqlandi. Shuning uchun ham dalachoy o‘simligidan foydalanib, sintetik preparatlardan farqli ravishda yon ta’sirlari kam yangi oziq-ovqat qo‘shilmasini yaratish va qandli diabet hamda yurak-qon tomir kasalliklarida qo‘llash maqsadga muvofiq degan xulosaga keldik.

**Kalit so‘zlar:** Endotelial, qon-tomir, antioksidant, immun, funksiyasi, ichak, rodentium, flavonoid, rutin, glibenklamid, qandli diabet.

**Abstract:** Studying the composition of the valerian plant revealed that it is rich in flavonoids. Therefore, we came to the conclusion that it is appropriate to create a new food additive with fewer side effects using the field plant and to use it in diabetes and cardiovascular diseases.

**Key words:** Endothelial, vascular, antioxidant, immune, function, intestine, rodentium, flavonoid, rutin, glibenclamide, diabetes.

**Аннотация:** Изучение состава растения валерианы выявило, что оно богато флавоноидами. Поэтому мы пришли к выводу, что на основе полевого растения целесообразно создать новую пищевую добавку с меньшим количеством побочных эффектов и использовать ее при диабете и сердечно-сосудистых заболеваниях.

**Ключевые слова:** эндотелиальная, сосудистая, антиоксидантная, иммунная, функция, кишечник, грызуны, флавоноиды, rutin, гlibenklamid, сахарный диабет.

**Kirish.** Flavonoidlar o‘simliklar tarkibida xalq tabobatida turli kasalliklar uchun tabiiy davolovchi vositalar sifatida ishlatiladi. Ularning juda ko‘p tajribalar orqali antioksidant xususiyatlari isbotlangan va inson salomatligi uchun foydali ekanligi ko‘rsatib berilgan hamda, tabiiy mahsulotlar neyroprotektor sifatidagi ta’siri tufayli ko‘proq e’tiborga sazovor bo‘lgan, chunki ular tabiatan bitta yo‘nalishda emas, balki pleyotrop ta’sir ko‘rsatadi[23;19].

Kversetinni og‘iz orqali 150-730mg/sutka dozada 4 haftadan 10haftagacha qabul qilish antigipertenziv ta’sir etishi aniqlangan[1]. Kversetin mRNK ekspresiyasini kamaytirish orqali renin-angiotenzin sistemasini ishini to‘xtatadi va shu bilan qon hajmi ortishini oldini oladi[2]. Qonda TAG lar miqdorini kamatirib, ZYULP tarkibida xolesterin miqdorini oshiradi. NO hosil bo‘lishini tezlashtirib endotelial vazodilatatsiyani ta‘minlaydi. Endotelial hujayralar apoptozini kamaytiradi[3;4;5]. Qon-tomir sistemasini Kversetin hayvonlar tanasiga yuborilganda antioksidant va immun tizim faolligi oshganini kuzatishgan[6]. Turli o‘tkazilgan tajribalar oksidlangan yog‘ ta’sirida ichaklardagi oksidlanishli stress kversetin ta’sirida pasaygani va oksidlanishli qaytarilish balansi yaxshilanib, ichak morfologik

strukturasi va baryer funksiyasi tiklangani kuzatilgan[7].Kversetin ichak mikroflorasi balansini ta'minlaydi.Bu modda ta'sirida ichak mikrobiotasini modulyatsiyalanishi kuzatilgan. Kversetin *C. rodentium* bilan bog'liq bo'lgan kolitni yengillashtiradi va yallig'lanishni kamaytiradi[8].

Hozirgacha diabetni to'liq davolovchi dorilar mavjud emas. Diabetda ishlatilayotgan dorilarning ayrimlari yon ta'sirlarga ega yoki uzoq qo'llanishi natijasida o'z ta'sirini yo'qotadi. Shuning uchun ham iloji boricha tabiiy vositalardan foydalanib bu kasallikni davolash maqsadga muvofiq. Chunki tabiiy vositalar yon ta'sirga ega emas. Tabiiy flavonoid rutin esa  $\alpha$  glyukozydazaga ta'sir etishi ma'lum va shu xususiyati tufayli qandli diabetda qo'llanilishi mumkun[9]. Bu flavonoid boshqa flavonoidlar bilan taqqoslanganda qandli diabetlarda glyukoza tolerantlikni oshirishi, FBG va lipidlar miqdorini kamaytirishi isbotlangan. Rutinni ta'siri glibenklamid bilan taqqoslanganda diabet bilan kasallangan kalamushlarda glyukoza miqdorini pasaytirishini aniqlashgan[10]. Shrinivasan va boshqalar streptozototsin bilan qandli diabet chaqirilgan kalamushlarda rutin muntazam qabul qilish orqali 30-40% giperqlikemiyani oldi olinganini ko'rsatishgan[11].

Rutin diabetlarda xolesterinni so'rilishini ingibirlaydi. Qandli diabeti bor bemorlarda qonda ZPLP miqdori kutarilib, ZYULP miqdori kamayib ketadi. Ularni statinlar va fibratlar kabi gipolipidemik preparatlar bilan davolash ZPLP miqdorini  $<70$  mg/dl gacha tushurmaydi[12]. O'tkazilgan klinik tajribalar dorivor o'simliklar tarkibida mavjud bo'lgan rutin qonda lipidlar miqdoriga ijobiy ta'sir etishini ko'rsatdi[13;14]. Adabiyotlarda rutin saqlovchi dorivor o'simliklar berilganda qonda ZPLP, ZJPLP va TAG lar miqdorining sezilarli darajada pasayishini alohida ta'kidlangan[15;16;10;18]. Rutin ta'sirida ZYULP miqdorini oshishi ham qator eksperimental va klinik tajribalarda isbotlangan[19;20;21].

Digidrokversetinni 21 kun davomida kalamushlarga berilganda xolesterin miqdorini kamayishi kuzatilgan. Chunki bu flavonoid GMG-KoA reduktaza ingibitori bulib, organizmda xususan jigarda xolesterin sintezini ingibirlaydi.Xolesteringa boy ozuqa bilan boqilgan kalamushlarga digidrokversetin berilganda axlatida xolesterin miqdori oshgani kuzatilgan[23]. AroA-I va ApoB lar ZPLP va ZYULP larning retseptorlariga ligand sifatida ta'sir etadi. ApoB to'qimalarni xolesterin bilan boyitsa, AroA-I esa to'qimalardan jigarga xolesterin tashilishini kuchaytiradai. Digidrokversetin ta'sirida AroA-I miqdori oshib, ApoB miqdori esa kamayishi kuzatildi.Bu esa digidrokversetinga boy o'simliklar bilan ratsionni boyitish orqali giperlipidemiyani oldini olish mumkunligini ko'rsatdi[24].

Digidrokversetin gepatoprotektorlik xususiyatiga ham ega bo'lib, digidrokversetin ta'sirida alaninaminotransferaza va aspartaminotransferazalarning miqdori sezilarli kamaygan. Konkanavalin A (Con A) qabul qilgan kalamushlarda digidrokversetin immunoregulyatorlik funksiyalari tekshirilgan va ijobiy natijalar olingan. Masalan: HepG2 hujayralarida TNF- $\alpha$ /ActD indutsirlagan apoptozdan kaspaza-3, kaspaza-7 va kaspaza-8 aktivligini ingibirlash orqali himoya qilgan. Bundan tashqari digidrokversetin NF-kB/p65ni fosforillanishini kamaytirib, proapoptik faktorlarni ekspressiyasini ingibirlaydi va antiapoptik faktorlardan biri bo'lgan Bcl-2 ni ekspressiyasini kuchaytirgan. Jigarning T- hujayralari CD4 + i CD8 + infiltratsiyasi va yallig'lanish mediatorlari ekspressiyasini kamaytirish orqali jigar hujayralarining immunologik yemirilishlarini kamaytirgan. Digidrokversetin HepG2 hujayralarini TNF- $\alpha$  dan himoyalagan[22].

Lyuteonin LXR $\alpha$  glyukoneogenez fermentlari aktivligini pasaytiradi va glyukoza shzlashtirilishini yaxshilaydi. xolesterin-7- $\alpha$ -gidroksilaza, ABCG1 va SRB1 ekspressiyasini

kuchaytiradi[25]. Lyuteonin va gall kislotasi og‘ir metallar bilan zaharlanishni oldini oladi. Masalan ayrim kosmetik vositalar va lak-bo‘yoqlar tarkibida qo‘rg‘oshin bulib, uning organizmga tushishi salbiy oqibatlarga olib kelishi mumkun[26]. Ayrim soha vakillari tomonidan kobalt saqllovchi ozuqa va ichimliklarni iste‘mol qilish, sportchilar tomonidan taqiqlangan doping qabul qilish va ortopedik protezlar orqali kobalt organizmga tushishi mumkun. Bu esa kognitiv funksiyalar va kordinatsiyaning buzilishi hamda depressiyaga olib kelishi mumkun[27].

Adabiyotlar tahlilidan ko‘rinib turibdiki, flavonoidlar qon-tomir kasalliklarida gipolipedemik ta‘sir ko‘rsatadi, qandli diabetda esa glyukoneogenezni ingibirlaydi hamda glyukozani hujayralarga o‘tishini yaxshilaydi. Undan tashqari gepatoprotektor,immunoregulyatorlik funksiyalari mavjud.


**Tadqiqotning maqsadi.** Dalachoy tarkibidagi flavonoidlarni suyuqlik xromatografiyasi usuli yordamida aniqlash.

**Tadqiqot metodologiyasi.** Namuna tarkibidagi flovonoidlarni suyuqlik xromatografiyasi usuli yordamida aniqlandi. Namunadan 5-10 gr miqdorida analitik tarozida tortib olinib, 300 ml hajmdagi yassi kolbaga solinadi. Ustiga 50 ml 70% li etanol eritmasidan qo‘shildi. Aralashma magnit aralashtirgich, teskari sovutkich bilan jihozlanib, 1 soat davomida intensiv aralashtirib turgan holda 70-800S qaynatildi va keyinchalik 2 soat davomida xona haroratida aralashtirildi. Aralashma tindirilib filtrlab olindi. Qolgan qismiga 25 ml 70 % etanoldan solib 2 marta qayta ekstraksiya qilinadi. Filtratlar birlashtiriladi va 100 ml o‘lchagich kolbaga solinib chizig‘igacha 70% etanol bilan to‘ldirildi. Hosil bo‘lgan eritma sentrafugada 6000-8000 oborot/minut tezlikda 20-30 daqiqa davomida aylantiriladi. Hosil bo‘lgan eritma ustki qismidan analiz uchun olindi.


Adabiyotlarda Flovonoidlarni YUSSX bilan aniqlashda elyuyent sifatida fosforli, atsetatli bufer sistemalari va atsetonitrildan foydalanilgan. Biz fosfatli bufer sistemasi hamda atsetonitrildan foydalandik.

VEJX «Agilent-1200» markali Kolonka Agilent C18 5mkm,4,6x250mm. Elyuirovalanish izokratik rejimda o‘tkazildi, Mobil faza sifatida ),1%li ortofosfat kislotasi va atsetonitril smesi 70:30 nisbatda ishlatildi. Elyuyent potogi obyemli tezligi– 1,0 ml/min, kiritilgan namuna hajmi 10mkl. Uzunligi 254 nm.

Xromotografga dastlab, ishchi standart eritmalar, keyinchalik tayyorlangan ishchi eritmalar kiritildi.



Dalachoy standart eritmasi xromatogrammasi



Dalachoy ishchi eritmasi xromatogrammasi

Dalachoy o‘simligi tarkibidagi flavonoidlar.

	Dala choy
	Konsentratsiya mg/gr
Digidrokversitin	5,36
Lyutionin	0,51
Rutin	2,31
Kversitin	9,21
Gall kislotasi	0,016

Dalachoy o‘simligi tarkibi yuqori samarali suyuqlik xromatografiyasida o‘rganilganda, flavonoidlardan digidrokversetin, lyutionin, rutin, kversetin va gall kislotasi aniqlandi. Dalachoy tarkibida flavonoidlardan kversetin miqdori nisbatan ko‘pligini ko‘rdik. Digidrokversetin 5,36 mg/gr, lyutionin 0,51mg/gr, rutin 2,31mg/gr, kversetin 9,21 mg/gr ni tashkil qilishi aniqlandi.

XULOSA. Dalachoy o‘simligi tarkibi o‘rganilganda flavonoidlarga boy ekani aniqlandi. Shuning uchun ham dalachoy o‘simligidan foydalanib, sintetik preparatlardan farqli ravishda yon ta’sirlari kam yangi oziq-ovqat qo‘shilmasini yaratish va qandli diabet hamda yurak-qon tomir kasalliklarida qo‘llash maqsadga muvofiq degan xulosaga keldik.

### Foydalanilgan adabiyotlar.

1. Edwards R.L., Lyon T., Litwin S.E., Rabovsky A., Symons J.D., Jalili T. Quercetin reduces blood pressure in hypertensive subjects. *J. Nutr.* 2007;137:2405–2411. [PubMed] [Google Scholar]
2. Olaleye M.T., Crown O.O., Akinmoladun A.C., Akindahunsi A.A. Rutin and quercetin show greater efficacy than nifedipin in ameliorating hemodynamic, redox, and metabolite imbalances in sodium chloride-induced hypertensive rats. *Hum. Exp. Toxicol.* 2014;33:602–608. doi: 10.1177/0960327113504790. [PubMed] [CrossRef] [Google Scholar]
3. Pfeuffer M., Auinger A., Bley U., Kraus-Stojanowic I., Laue C., Winkler P., Rufer C.E., Frank J., Bosch-Saadatmandi C., Rimbach G., et al. Effect of quercetin on traits of the metabolic syndrome, endothelial function and inflammation in men with different APOE isoforms. *Nutr. Metab. Cardiovasc. Dis.* 2013;23:403–409. doi: 10.1016/j.numecd.2011.08.010. [PubMed] [CrossRef] [Google Scholar]
4. Erdogan A., Most A.K., Wienecke B., Fehsecke A., Leckband C., Voss R., Grebe M.T., Tillmanns H., Schaefer C.A., Kuhlmann C.R. Apigenin-induced nitric oxide production involves calcium-activated potassium channels and is responsible for antiangiogenic effects. *J.*

Thromb. Haemost. 2007;5:1774–1781. doi: 10.1111/j.1538-7836.2007.02615.x. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

5. Dayoub O., Andriantsitohaina R., Clere N. Pleiotropic beneficial effects of epigallocatechin gallate, quercetin and delphinidin on cardiovascular diseases associated with endothelial dysfunction. *Cardiovasc. Hematol. Agents Med. Chem.* 2013;11:249–264. doi: 10.2174/1871525712666140309233048. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

6. Song ZhangabIn, HoKim. Effect of quercetin (flavonoid) supplementation on growth performance, meat stability, and immunological response in broiler chickens. *Livestock Science*. Volume 242, December 2020, 104286

7. Yuanyang Dong, Jiaqi Lei, Bingkun Zhang. Effects of dietary quercetin on the antioxidative status and cecal microbiota in broiler chickens fed with oxidized oil. *Poult Sci.* 2020 Oct; 99(10): 4892–4903.

8. Rui Lin, Meiyu Piao, Yan Song. Dietary Quercetin Increases Colonic Microbial Diversity and Attenuates Colitis Severity in Citrobacter rodentium-Infected Mice. *Front Microbiol.* 2019 May 16;10:1092. doi: 10.3389/fmicb.2019.01092.

9. Solomon Habtemariam 1, Giovanni Lentini. The therapeutic potential of rutin for diabetes: an update. *Mini Rev Med Chem.* 2015;15(7):524-8. doi: 10.2174/138955751507150424103721.

10. R. Jadhav, G. Puchchakayala, Hypoglycemic and antidiabetic activity of flavonoids: boswellic acid, ellagic acid, quercetin, rutin on streptozotocin-nicotinamide induced type 2 diabetic rats, *Int. J. Pharm. Pharm. Sci.* 4 (2) (2012) 251–256.

11. K. Srinivasan, C. Kaul, P. Ramarao, Partial protective effect of rutin on multiple low dose streptozotocin-induced diabetes in mice, *Ind. J. Pharmacol.* 37 (5) (2005) 327

12. A.W. Dake, N.D. Sora, Diabetic dyslipidemia review: an update on current concepts and management guidelines of diabetic dyslipidemia, *Am. J. Med. Sci.* 351 (4) (2016) 361–365.

13. A. Ghorbani, Clinical and experimental studies on polyherbal formulations for diabetes: current status and future prospective, *J. Integr. Med.* 12 (4) (2014) 336–345.

14. A. Ghorbani, Phytotherapy for diabetic dyslipidemia: evidence from clinical trials, *Clin. Lipidol.* 8 (3) (2013) 311–319

15. P. Prince, N. Kannan, Protective effect of rutin on lipids, lipoproteins, lipid metabolizing enzymes and glycoproteins in streptozotocin-induced diabetic rats, *J. Pharm. Pharmacol.* 58 (10) (2006) 1373–1383.

16. Y.B. Wang, Z.M. Ge, W.Q. Kang, Z.X. Lian, J. Yao, C.Y. Zhou, Rutin alleviates diabetic cardiomyopathy in a rat model of type 2 diabetes, *Exp. Ther. Med.* 9 (2) (2015) 451–455.

17. O.M. Ahmed, A.A. Moneim, I.A. Yazid, A.M. Mahmoud, Antihyperglycemic, antihyperlipidemic and antioxidant effects and the probable mechanisms of action of Ruta graveolens infusion and rutin in nicotinamide-streptozotocin-induced diabetic rats, *Diabetol. Croat.* 39 (1) (2010) 15–35.

18. Hawthorn flavonoid ameliorates cognitive deficit in mice with Alzheimer's disease by increasing the levels of Bifidobacteriales in gut microbiota and docosapentaenoic acid in serum metabolites. *Food Function.* 2022 Nov 28;13(23):12371-12382. doi: 10.1039/d2fo02871a.

19.A.A.H. Fernandes, E.L.B. Novelli, K. Okoshi, M.P. Okoshi, B.P. Di Muzio, J.F.C. Guimarães, A.F. Junior, Influence of rutin treatment on biochemical alterations in experimental diabetes, *Biomed. Pharmacother.* 64 (3) (2010) 214–219

20.H.R. Sadeghnia, B.S. Yousefsani, M. Rashidfar, M.T. Boroushaki, E. Asadpour, A. Ghorbani, Protective effect of rutin on hexachlorobutadiene-induced nephrotoxicity, *Renal Fail.* 35 (8) (2013) 1151–1155.

21. Jiajie Chen, Xu Sun, Tingting Xia, Qiqi Mao, Liang Zhong. Pretreatment with dihydroquercetin, a dietary flavonoid, protected against concanavalin A-induced immunological hepatic injury in mice and TNF- $\alpha$ /ActD-induced apoptosis in HepG2 cells. *Food Function.* 2018 Apr 25;9(4):2341-2352.

doi: 10.1039/c7fo01073g.

22. Franziska Pohl 1, Paul Kong Thoo Lin. The Potential Use of Plant Natural Products and Plant Extracts with Antioxidant Properties for the Prevention/Treatment of Neurodegenerative Diseases: In Vitro, In Vivo and Clinical Trials. *Molecules.* 2018 Dec 11;23(12):3283. doi: 10.3390/molecules23123283.

23. Itaya, S., Igarashi, K., 1992. Effects of taxifolin on the serum cholesterol level in rats. *Biosci. Biotechnol. Biochem.* 56, 1492–1494.

24. Igarashi, K., Uchida, Y., Murakami, N., Mizutani, K., Masuda, H., 1996. Effect of a chrysolepis on the serum and liver lipid concentrations and on the erythrocyte and liver antioxidative enzyme activities of rats. *Biosci. Biotechnol. Biochem.* 60, 513–515.

25. Hee-Sook Park, Kyunhee Lee, Soon-Hee Kim, Moon Ju Hong, Nam-Joo Jeong, Myung-Sunny Kim. Luteolin improves hypercholesterolemia and glucose intolerance through LXR $\alpha$ -dependent pathway in diet-induced obese mice. *Journal of Food Biochemistry.* Volume 44, Issue 9 e13358 <https://doi.org/10.1111/jfbc.13358>

26. Baty R. S., Hassan K. E., Alsharif K. F., et al. Neuroprotective role of luteolin against lead acetate-induced cortical damage in rats. *Human & experimental toxicology* . 2020;39(9):1200–1212. doi: 10.1177/0960327120913094. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

27. Akinrinde A. S., Adebisi O. E. Neuroprotection by luteolin and gallic acid against cobalt chloride-induced behavioural, morphological and neurochemical alterations in Wistar rats. *Neurotoxicology* . 2019;74:252–263. doi: 10.1016/j.neuro.2019.07.005. [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]