

OXIDATION OF DIFFERENT SUBSTANCES AND Mg^{2+} - ACTIVATED ATPase IN MITOCHONDRIA OF BROWN ADIPOSE TISSUE

Turayeva Fatima Rajaboyevna¹
Mirzaolimov Mirzahid Mirzavaliyevich²

¹Namdu is a PhD candidate

²PhD, Namangan State University

E-mail: turayevafotima1612@gmail.com

E-mail: mirzohid_0421@mail.ru

Abstract: In mitochondria of brown adipose tissue, it has been studied the oxidation of various substrates, including NADH and ascorbate + cytochrome *c*, and determined the activation of Mg^{2+} -stimulated ATPase. It was revealed that NAD-dependent substrates (pyruvate + malate, glutamate) are oxidized with a high level of V_4 and low RC. However, on such substrate as succinate, it was detected a higher V_4 rate and a lower RC than the oxidation of NAD-dependent substrate. As well, high oxidation rates of exogenous NADH and ascorbate + cytochrome *c* were detected, not coupled with the ATP synthesis. At application of GDP, it was found a decrease in the respiration rate of V_4 on succinate, on NAD-dependent substrates as well as on the NADH substrate.

At the same time, liver mitochondria have been studied, that have mildly expressed uncoupled respiration at oxidation of NADH, ascorbate + cytochrome *c* and other substrates.

In mitochondria of adipose tissue, it was found a high activity of Mg^{2+} -stimulated ATPase, sensitive to oligomycin. It has a higher activity than DNP-stimulated ATPase.

In liver mitochondria, it is detected a significantly lower level of exogenous NADH oxidation as well as Mg^{2+} -stimulated ATPase than in mitochondria of brown adipose tissue.

The intensive oxidation of exogenous NADH and ascorbate (+ cytochrome *c*), as well as high activity of Mg^{2+} -stimulated ATPase, detected in mitochondria of brown adipose tissue are explained by the presence of a special membrane-porous subpopulation of mitochondria in suspension. This subpopulation, that exists separately from the ATP-synthesizing population of mitochondria, is capable of carrying out intense proton leakage and tissue thermogenesis.

Keywords: mitochondria of brown adipose tissue, proton leakage, coupled and uncoupled respiration, thermogenesis, bioenergetics, UCP1.

Annotatsiya: yog' to'qimasi, u turli substratlarning, shu jumladan NADH va askorbat + sitoxrom *c* oksidlanishini o'rganib chiqdi va Mg^{2+} bilan stimulyatsiya qilingan ATPazning faollashishini aniqladi. NADga bog'liq substratlar (piruvat + malat, glutamat) yuqori darajadagi V_4 va past RC bilan oksidlanishi aniqlandi. Biroq, süksinat kabi substratda NADga bog'liq substratning oksidlanishiga qaraganda yuqori V_4 darajasi va pastroq RC aniqlandi. Shuningdek, ekzogen NADH va askorbat + sitoxrom *c* ning yuqori oksidlanish tezligi, ATP sintezi bilan birlashtirilmaganligi aniqlandi. Ya'ni qo'llashda süksinatda, NADga bog'liq substratlarda, shuningdek NADH substratida V_4 ning nafas olish tezligining pasayishi aniqlandi.

Shu bilan birga, jigar mitoxondriyalari o'rganildi, ular NADH, askorbat + sitoxrom *c* va boshqa substratlarning oksidlanishida engil ifodalanmagan nafas olishni ifodalaydi.

Yog' to'qimalarining mitoxondriyalarida oligomisinga sezgir bo'lgan Mg^{2+} -stimulyatsiya qilingan ATPazning yuqori faolligi aniqlandi. U DNP-stimulyatsiya qilingan ATPazaga qaraganda yuqori faollikka ega.

Jigar mitoxondriyalarida jigarrang yog 'to'qimalarining mitoxondriyalariga qaraganda ekzogen NADH oksidlanish darajasi, shuningdek Mg^{2+} -stimulyatsiya qilingan ATPazning sezilarli darajada pastligi aniqlanadi.

Jigarrang yog' to'qimalarining mitoxondriyalarida aniqlangan ekzogen NADH va askorbat (+ sitoxrom c) ning intensiv oksidlanishi, shuningdek, Mg^{2+} -stimulyatsiyalangan ATPazaning yuqori faolligi suspenziyada mitoxondriyaning maxsus membrana-g'ovakli subpopulyatsiyasi mavjudligi bilan izohlanadi. Mitoxondriyaning ATP sintez qiluvchi populyatsiyasidan alohida mavjud bo'lgan bu subpopulyatsiya intensiv proton oqishini va to'qimalarning termogenezini amalga oshirishga qodir.

Kalit so'zlar: jigarrang yog 'to'qimalarining mitoxondriyalari, proton oqishi, qo'shilgan va qo'shilmagan nafas olish, termogenez, bioenergetika, UCP1.

Абстрактный: жировой ткани, изучено окисление различных субстратов, в том числе НАДН и аскорбата + цитохрома с, и определена активация Mg^{2+} -стимулируемой АТФазы. Выявлено, что НАД-зависимые субстраты (пируват+малат, глутамат) окисляются с высоким уровнем V4 и низким RC. Однако на таком субстрате, как сукцинат, обнаружена более высокая скорость V4 и более низкий RC, чем при окислении НАД-зависимого субстрата. Также обнаружены высокие скорости окисления экзогенного НАДН и аскорбата + цитохрома с, не сопряженного с синтезом АТФ. При применении ВВП обнаружено снижение скорости дыхания V4 на сукцинате, на НАД-зависимых субстратах, а также на субстрате НАДН.

В то же время изучены митохондрии печени, имеющие умеренно выраженное несвязанное дыхание при окислении НАДН, аскорбата + цитохрома с и других субстратов.

В митохондриях жировой ткани обнаружена высокая активность Mg^{2+} -стимулируемой АТФазы, чувствительной к олигомицину. Он имеет более высокую активность, чем DNP-стимулируемая АТФаза.

В митохондриях печени обнаруживается достоверно меньший уровень экзогенного окисления НАДН, а также Mg^{2+} -стимулированной АТФазы, чем в митохондриях бурой жировой ткани.

Интенсивное окисление экзогенного НАДН и аскорбата (+цитохром с), а также высокая активность Mg^{2+} -стимулируемой АТФазы, обнаруженные в митохондриях бурой жировой ткани, объясняются наличием в суспензии особой мембранопористой субпопуляции митохондрий. Эта субпопуляция, существующая отдельно от АТФ-синтезирующей популяции митохондрий, способна осуществлять интенсивную утечку протонов и тканевой термогенез.

Ключевые слова: митохондрии бурой жировой ткани, утечка протонов, связанное и несвязанное дыхание, термогенез, биоэнергетика, UCP1.

Abbreviations: BAT - brown adipose tissue, ADP- adenosine diphosphate, ATP- adenosine triphosphate, GDP- guanosine diphosphate, NADH-nicotinamide adenine dinucleotide reduced, 2,4-DNP-2,4-dinitrophenol, RC – respiratory control.

1. Introduction.

The study of BAT of mitochondria was mainly of interest in clarifying the mechanism of biological thermogenesis (Brand, 1990; Ricquier and Bouillaud, 2000; Porter, 2001; Brand et al., 2005; Jastroch et al. 2010; Bal 2017).

Mitochondria, isolated from BAT usually have a high rate of V_4 respiration and weak coupling with ATP synthesis. This phenomenon has served as the main prerequisite for the promotion of the thermogenic uncoupling respiration idea in mitochondria of BAT (Ricquier, 2017). Subsequently it was established that the cause of proton leakage, uncoupling in mitochondria is the presence of a particular protein - thermogenin (UCP1) in their inner membrane (Ricquier, 2017). The isolated protein UCP1 easily build into liposomes and bilipid artificial membranes, that leads to an increase in their membrane permeability to proton (Brown, Brand, 1991; Ricquier, 2017) and activated by fatty acids and suppressed by GDP (Garlin, Jaburek, Jeiek, 1998; Fedorenko, Lishko, Kirichok, 2012). Increased proton permeability of mitochondria by UCP1 is regarded as the main reason of proton-driving power (Δp) directly into heat (Brand, 1990; Ricquier and Bouillaud, 2000; Porter, 2001; Brand, Pakav et al, 2005.). According to the literature (Hulbert, Else, Manolis, Brand, 2002; Jastroch, Divakaruni, et al, 2010) all mitochondria of BAT take part in proton leak. It is also shown that the proton leakage is active also in the conditions of the brown adipose tissue cell (Silva, Oliveira, 2012; Mahdaviyani, Benador, Shirihai, 2015).

In literature (Brand, 1990; Hulbert, 2002; Bal, Singh, 2017), at study of proton leakage in mitochondria of BAT and other tissues, succinate is used as a main oxidation substrate. In a number of studies (Cannon, Nedergaard, 2004; Nicholls, 2006; Shabalina et al., 2010; De Meis et al., 2012), it was made a comparison of proton leakage intensity with the use of pyruvate + malate and succinate as oxidation substrates in mitochondria. In these studies, oxidation of succinate occurs with less coupling than in the oxidation of pyruvate + malate. Therefore, it is noted (De Meis, Ketzer et al., 2012) that succinate can be oxidized in a different way than NAD-dependent substrates. We carried out such an analysis earlier (Akhmerov, 1981; Akhmerov, 1986). According to our data, this way may be an uncoupling system of mitochondria that oxidizes not only succinate, but also NADH, ascorbate (+cytochrome *c*). However, these reactions remained poorly studied at BAT mitochondria (Akhmerov, Makhmudov, 1988) that were the prerequisite for carrying out this experimental work.

2. Materials and methods

Adult rats were used for experiments. The sacrifice of animals was carried out by decapitation. Interscapular part of the animals' body was opened quickly by the scissors and BAT was extracted. The tissue was immediately immersed in a cooled solution of isolation medium and then its mass was determined.

2.1. Isolation of mitochondria from BAT and liver

To obtain mitochondria, we were guided by a method adapted for this tissue (Grav, Pederson, Christiansen, 1970; Skaane, Christiansen et al., 1972). The basic incubation medium used by us contained 300 mM sucrose, 0.5% bovine serum albumin (fat-free BSA), 10 mM Tris-HCl (pH 7.5). To obtain of BAT mitochondria, the raised amount of fatty acids in this tissue was taken into account. Therefore, as in some works (Grav, Pederson, 1970; Skaane, Christiansen et al., 1972), we used 0.5% albumin (defatted) in the composition of incubation medium. For the grinding of the tissues, a glass homogenizer with a Teflon pestle was used. Homogenization of the tissue was carried out in a 10-fold volume of mentioned

incubation medium for 20-30 seconds. The obtained homogenate was centrifuged at 700 g for 7 minutes to precipitate large cell aggregates. Mitochondria were precipitated from the supernatant by repeated centrifugation at $6000 \times g$ for 20 minutes. The supernatant was removed and the mitochondrial pellet was again diluted in a 5-fold volume of incubation medium and centrifuged at $6000 \times g$ for 20 minutes. The pellet was suspended in 0.1 ml of incubation medium to the initial tissue weight.

In a comparative plan, it was of interest to isolate rat liver mitochondria under the same conditions as BAT mitochondria. Therefore, a part of the experiments was performed on liver mitochondria with the use of different oxidation substrates.

All procedures for obtaining of mitochondrial preparations were performed at temperatures of $-2 + 2^{\circ}\text{C}$.

2.2. Polarographic study of mitochondrial respiration

In the obtained mitochondrial preparations, phosphorylation and uncoupled respiration were determined by the generally accepted polarographic method (Chance, Williams, 1955). The polarographic cell had a volume of 1 ml. The incubation medium for performing the basic polarographic studies contained 120 mM KCl, 2 mM EGTA, 5 mM KH_2PO_4 , 10 mM tris-HCl (pH 7.5). The used oxidation substrates are indicated in the tables.

The amount of BAT mitochondria, inserted into the polarographic cell was about 2 mg/ml. The protein was determined by Lowry et al. (Lowry, Rosebrough et al., 1951).

The respiration rate of the preparations is expressed in nanograms of atoms of absorbed oxygen per minute per mg of protein (ng-atom O/min per mg of mitochondrial protein).

In separate experiments on BAT mitochondria, it was tested the effect of GDP (0.5 mM aqueous solution, pH 7.5) to the oxidation of the used substrates.

2.3. Determination of ATPase activity

Determination of ATPase activity of mitochondria was carried out pH-metrically (Chance, Nishimura, 1967) with registration on the recorder. The main incubation medium consisted of 120 mM KCl and 2 mM tris-HCl (pH 7.5). In the cell, either BAT mitochondria (about 2 mg of mitochondrial protein) or liver mitochondria (about 5 mg mitochondrial protein) was added.

In the study of Mg^{2+} -dependent ATPase, mitochondria were first added to the measuring cell, then 1 mM of ATP. Activation of ATPase was caused by the addition of 3 mM MgCl_2 to the measuring cell.

At study of DNP-stimulated ATPase, incubation medium with 1 mM of ATP was used in experiments, as well as 2 mM of EDTA to bind bivalent cations. Under these conditions, the activation of ATPase was induced by the addition of 50 μM DNP.

To analyze of ATPase reactions following substances were also used: inhibitors – oligomycin 1.5 $\mu\text{g}/\text{ml}$ and carboxyatractyloside 1 $\mu\text{g}/\text{ml}$. The value of ATPase activity was expressed in $\mu\text{M H}^+/\text{min}$. per mg of mitochondrial protein. The temperature of the preparations' incubation is 25°C .

3. Results and discussion

In order to ascertain the energy metabolism features in BAT mitochondria, we studied the intensity of oxidation of different substrates in isolated mitochondria of this tissue in adult rats.

3.1. Endogenous respiration of BAT mitochondria

First, it was interesting to study the endogenous respiration of mitochondria, that is, without the addition of exogenous substrates. The level of this respiration was high and it did not react to ADP additives, which indicates the uncoupling of oxidative phosphorylation in mitochondria (Table 1, Fig. 1). The reason for the uncoupling may be fatty acids content in mitochondrial suspension (Nicholls, 2006; De Meis., Ketzer, 2012), that can cause an increase in endogenous respiration. At first, proton leakage occurs through UCP1 (Brand, Pakav et al., 2005; Jastroch, Divakaruni et al., 2010; Bal, Singh, 2017), then mitochondrial respiration uncoupled by analogy to 2,4-DNP, and mitochondrial respiration may also increase as a result of the oxidation of acyl derivatives of fatty acids (Nicholls, 2006). Even a single wash of mitochondria with addition of purified BSA slightly reduces endogenous respiration. However, the introduction of BSA into the incubation medium significantly reduces the endogenous respiration of mitochondria (Akhmerov, Makhmudov, 1988; Nicholls, 2006), possibly due to the binding of fatty acids.

3.2. Oxidation of different substrates

We clarified the features of different substrates oxidation in BAT mitochondria, in particular, NAD-dependent substrates and succinate. The obtained results are shown in Table 1 and Fig. 1.

Table 1. Oxidation of different substrates in BAT mitochondria

Oxidation substrates	V ₃	V ₄	RC
pyruvate + malate	64.17±5.43	38.21±5.14	1.68
Succinate	106.51±6.42	98.63±6.35	1.18
NADH	126.75±8.73	126.75±8.73	1.0
NADH+GDP	77,45 ± 6,14	77,45 ± 6,14	1.0
NADH + rotenone	3.8±0.45	3.8±0.45	1.0
ascorbate + cytochrome <i>c</i>	211.14±13.14	211.14±13.14	1.0
ascorbate + cytochrome <i>c</i> + sodium cyanide	14.12±1.16	14.12±1.16	1.0

Note: In this table, rates of mitochondrial respiration (V₃, V₄,) are given in ng atom of O/min. per mg of protein. Concentration of GDP 0.5 mM, NADH 1 mM. At use of ascorbate, 250 µg/ml of cytochrome C was additionally supplemented into incubation medium.




Fig. 1. Polarographic records of BAT mitochondrial respiration of newborn guinea pigs. Abbreviation: olig. – oligomycin, rot. - rotenone, cyt.*c* – cytochrome *c*, CN⁻ – sodium cyanide. The figures show the respiration rate. -GDP – respiration in the absence of GDP, +

GDP - respiration in the presence of GDP (1 mM). Substrates of oxidation are indicated above the polarograms. Mitochondrial protein is 1.4 mg per ml of incubation medium.

In BAT mitochondria, pyruvate + malate are oxidized with low coupling with phosphorylation, that is combined with a high rate of V_4 respiration. In this case, the reduction in the coupling parameters can result in endogenous fatty acids. At use of succinate as an oxidation substrate, an even lower level of coupling is revealed, along with an increase in V_4 respiratory rate, that additionally indicates the uncoupling of mitochondrial respiration. These results are in complete agreement with the literature data ([Brand et al., 2005](#); [Jastroch, Divakaruni et al., 2010](#); [De Meis et al., 2012](#)). Weak coupling is explained by the presence of mitochondrial uncoupling protein - UCP1 in the inner membrane, that activates proton leakage. Basically, this conclusion was made, with the use of succinate in mitochondria ([Brand, 1990](#); [Hulbert et al., 2002](#)). We used different oxidation substrates in experiments. The obtained data showed that uncoupling of mitochondrial respiration is not the same in the oxidation of different substrates. A more significant uncoupling of respiration occurs in succinate oxidation that is evident from the high level of V_4 respiration than in the oxidation of NAD-dependent substrates. The high V_4 rate decreases with the action of GDP (Fig. 1), that indicates its connection with the proton leakage. However, it is not suppressed by oligomycin, i.e. this respiration is not associated with the ATP synthesis and utilization.

These results motivated us to search for the cause of high uncoupling of mitochondrial respiration during succinate oxidation in comparison with NAD-dependent substrates. Earlier it was shown that on cardiac mitochondria this difference between substrates is more distinct, since they have low endogenous respiration ([Akhmerov, 1986](#), [Akhmerov, Niyazmetov, 2016](#)). By analogy with cardiac mitochondria, we assumed that some mitochondria function in an uncoupled mode in the suspension. The membrane of this group of mitochondria can be highly porous permeable to succinate, possibly for NADH, as well as ascorbate (+ cytochrome *c*). For us, the nature of oxidation of added NADH or ascorbate + cytochrome *c* of great interest.

Testing of NADH in BAT mitochondria gave unexpected results. It was oxidized with high intensity and its oxidation did not influence on added ADP or oligomycin (Fig. 1), but was strongly suppressed by the rotenone (a blocker of the first link of the mitochondrial respiratory chain). The use of ascorbate + cytochrome *c* also showed a high oxidation rate in mitochondria of BAT, which did not react to added ADP, and sodium cyanide (a blocker of the third link) dramatically suppressed the oxidation process (Fig. 1). Consequently, the last two substrates are oxidized along the main respiratory chain of mitochondria, and not coupled with phosphorylation process (without ATP synthesis). We would like to remind that the last two substrates are unpopular among researchers and usually they avoid their use in experiments. Oxidation of these substrates is associated with damage to mitochondria ([Lehninger, 1966](#)). This conclusion was made in the study of liver mitochondria, where exogenous NADH is oxidized very poorly. However, there are a number of reasons to consider their oxidation as a native feature of mitochondria. Usually, NADH does not pass through intact phosphorylating mitochondria ([Lehninger, 1966](#)), and its oxidation indicates the presence of mitochondria with a high membrane permeability. We do not exclude that UCP1 causes high permeability for NADH and ascorbate in mitochondria. So, earlier in the literature, intensive NADH oxidation in muscle mitochondria was shown, which was considered as a native process ([Blanchaer, 1964](#); [Rasmussen, 1969](#); [Rasmussen, Rasmussen, 1985](#)).

To interpret own data, we hypothesized that in suspension of BAT mitochondria as in suspension of heart mitochondria (Akhmerov, 1986; Akhmerov, Makhmudov, 1988), there are two types of mitochondria, where one type functions associated with ATP synthesis, another type - not coupled. According to this view succinate oxidation with high V_4 and low RC associated with its utilization simultaneously in both types (subpopulations) of mitochondria, uncoupled oxidation of NADH or ascorbate, in our opinion, is due to their utilization by only uncoupled mitochondria. NAD-dependent substrates are oxidized mainly in phosphorylating mitochondria, since they may not be oxidized in membrane-porous uncoupled mitochondria due to lack of coenzyme NAD^+ . The addition of NAD^+ into the incubation medium accelerates pyruvate + malate oxidation. Thus, the features of different substrates oxidation in BAT mitochondria can be explained by the presence of two types of mitochondria in suspension (Akhmerov, 1986).

To compare the nature of appearance of uncoupled reactions in BAT mitochondria, we also studied liver mitochondria. These mitochondria are often the subject of many studies, since they have good coupling at different substrates. In our studies, we used substrates that are oxidized uncoupling in BAT mitochondria. We would like to remind that liver mitochondria were isolated and incubated under the same conditions as BAT mitochondria. From the results of Table 2, it can be seen that NAD-dependent substrates (malate + pyruvate) have higher coupling rates than succinate, which corresponds to the literature data. In liver mitochondria, the respiratory rate V_4 is lower, and the degree of coupling of respiration exceeds the coupling of BAT mitochondria by 3-6 times.

Table 2. Oxidation of different substrates in liver mitochondria

Substrates	V_3	V_4	RC
pyruvate + malate	62.31±3.04	11.12±1,02	5.60
Succinate	76.23±5.75	19.12±1.21	3.96
NADH	17.24±1.44	17.24±1.44	1.0
ascorbate + cytochrome <i>c</i>	21.45±2.12	21.45±2.12	1.0




Fig. 2. pH-metric records of 2,4-DNP activated and Mg^{2+} dependent ATPase in BAT and liver mitochondria. Abbreviation: olig. – oligomycin 1.5 $\mu\text{g/ml}$, Cattr-

carboxyatractyloside 1 $\mu\text{g/ml}$, 2,4-DNP 50 μM . The protein BAT mitochondria is 2.3 mg/ml , the protein of liver mitochondria is 4.3 mg per ml of the incubation medium. Rich lines - 1 mM of EDTA was added, MgCl_2 additives - 3 mM . L- liver mitochondria, brown adipose tissue mitochondria – BAT.

In addition, the oxidation of added NADH along the rotenone-sensitive pathway in liver mitochondria occurs at a very low rate, which does not change from the subsequent ADP addition. We also revealed approximately the same low rate of oxidation on ascorbate + cytochrome *c*. Our data correspond to numerous literature on this profile. The well-known fact of inoxidizability of NADH by the rotenone-sensitive pathway is due to the poor permeability of phosphorylating liver mitochondria to these nucleotides (Lehninger, 1966). Therefore, uncoupled pathway of substrates oxidation in mitochondria is a poorly developed. However, in liver mitochondria generally there is a phosphorylating (ATP synthesizing) oxidation pathway. Most probably, in liver, the bulk of mitochondria consists of a coupled subpopulation. This circumstance can determine high rates of oxidative phosphorylation in NAD-dependent substrates and succinate. Perhaps, liver mitochondria have historically become the most popular subject of research because of the weak manifestation of uncoupled oxidation in them than in mitochondria of other tissues.

3.3. *Determination of Mg^{2+} -stimulated ATPase in BAT mitochondria*

The possibility of the presence of two types of mitochondria in a suspension of BAT is also shown at the study of ATPase activity. Experiments have shown that two types of mitochondria function in isolated mitochondria, one of which is Mg^{2+} -stimulated ATPase and is associated with ATP hydrolysis. It can be localized in membrane-porous mitochondria, since it is not sensitive to carboxyatractyloside. Another ATPase is 2,4-DNP-stimulated and is associated with phosphorylating mitochondria because it is sensitive to oligomycin.

It should be noted that Mg^{2+} -stimulated ATPase is approximately 5 times active in BAT mitochondria than in liver mitochondria. It is not inhibited by carboxyatractyloside, but inhibited by oligomycin very effectively.

Previously, Mg^{2+} -stimulated ATPase was described in muscles mitochondria (Chao, Devis, 1972) and was considered as a native component of mitochondria (Chao, Devis, 1972, Vijayakumar and Weidemann, 1976).

We compared the activity of Mg^{2+} -stimulated ATPase and DNP-stimulated ATPase in BAT mitochondria. As it can be seen from Fig. 2, the first ATPase shows a high activity and it is approximately 4 times higher than the activity of DNP-stimulated ATPase. From Fig. 2 it also can be seen that Mg^{2+} -stimulated ATPase, in contrast DNP-stimulated ATPase, is not suppressed by oligomycin. These data indicate the presence of two types of ATPases in BAT mitochondria, the first of which has high activity and does not participate in the ATP synthesis. The presence of two ATPases in suspension BAT mitochondria can be explained by the coexistence of two mitochondrial subpopulations, one of which synthesizes ATP and is well known among specialists of bioenergetics, and the other subpopulation carries out uncoupled oxidation of substrates and has Mg^{2+} stimulated ATPase.

Previously it has been shown repeatedly presence of Mg^{2+} stimulated ATPase activity in muscle (Chao, Davis, 1972; Saks, Chernousova, 1975) and spleen (Vijayakumar and Weidemann, 1976) mitochondria, which hydrolyzes ATP. The functional significance of this enzyme, as a whole, remains unknown, although the above authors consider it to be native. According to the available prerequisites creatine phosphokinase is directly related to this

issue. It is known that ATP, synthesized in mitochondria, immediately converted to creatine phosphate by means of creatine phosphate kinase. Sachs et al (Saks, Chernousova, 1975) consider that phosphocreatine is a transport form of macroergic phosphates, that may protect macroergs from Mg^{2+} stimulated ATPase activity. Therefore creatine phosphokinase protects ATP from the adverse effects of Mg^{2+} stimulated ATPase activity. In addition to this factor protein inhibitor of ATPase and ADP are known in the literature (Skulachev, 1989; Nakamura, Fujikawa, Yoshida; 2013; García-Bermúdez J. Cuezva J. M. 2016). According to current opinions last two agents at a low membrane potential of mitochondria (in uncoupled mitochondria) can act as effective regulators of Mg^{2+} stimulated ATPase (Skulachev, 1989). It has also been shown that BAT mitochondria contain an active creatine kinase system that can protect ATP from Mg^{2+} stimulated ATPase (Bertholet, et al., 2017).

A part of the experiments was performed with liver mitochondria on definition of DNP- and Mg^{2+} dependent ATPases. From Fig. 2 it can be seen that in this case mitochondria have a high DNP-stimulated ATPase, which is suppressed by the addition of oligomycin to the incubation medium. The determination of Mg^{2+} -stimulated ATPase of liver mitochondria showed, on the contrary, a low activity, which is 5-6 times lower than DNP-stimulated ATPase. The addition of oligomycin caused suppression of this activity that indicates mitochondrial origin of this ATPase. It can be assumed that its low activity in liver mitochondria is associated with a small amount of membrane-porous mitochondria in the cells of this tissue.

4. Conclusion

From the above data, it follows that BAT mitochondria are very different from liver mitochondria in terms of high levels of uncoupled respiration that is apparently due to the high level of membrane-porous mitochondrial populations in BAT cells.

As a result, it may be said that in BAT mitochondria there can be two subpopulations of mitochondria. One of them carries out phosphorylating respiration, and the other has a porous membrane and carries out intense uncoupled respiration, and possesses Mg -stimulated ATPase. This subpopulation of mitochondria is an integral part of proton leakage and tissue thermogenesis.

REFERENCES

1. Axmerov R. N. et al. ON THE POSSIBILITY OF UNCOUPLED MITOCHONDRIA IN BROWN FAT OF NEWBORN GUINEA PIGS //Scientific Bulletin of Namangan State University. – 2019. – T. 1. – №. 9. – C. 49-55.
2. Mirzaolimov M. M., Abdullaev G. R., Abdullayev S. S. ROLE OF A CALORIE-RESTRICTED DIET IN PROLONGING THE LIFESPAN OF AN ORGANISM AND ITS MITOCHONDRIAL MECHANISMS //Scientific Bulletin of Namangan State University. – 2019. – T. 1. – №. 10. – C. 106-112.
3. Niyazmetov B. et al. UNCOU'LED RESPIRATION IN BIRD MITICHONDRIA: CONNECTION WITH THERMOGENESIS //Scientific Bulletin of Namangan State University. – 2019. – T. 1. – №. 2. – C. 100-104.
4. Soliev N., Mirzaolimov M. ACTION OF CALCIUM ON THE CONTENT OF PHOSPHOTYDYLCHOLIN, PHOSPHATYL ETHANOLAMINE AND THEIR LYSOFORMS IN THE RAT LIVER MITOCHONDRIA //Scientific Bulletin of Namangan State University. – 2019. – T. 1. – №. 3. – C. 69-71.

5. Mirzaolimov M. M. et al. THE METHOD OF SEPARATION OF MITOCHONDRIAS AND DETERMINATION OF PHYSIOLOGICAL AND BIOCHEMICAL CHANGES IN ORGANISMS IN ONTOGENESIS //Scientific Bulletin of Namangan State University. – 2020. – Т. 2. – №. 3. – С. 175-178.
6. Мирзаолимов М. М., Рахимжонович М. А. Г. ВЛИЯНИЕ ПЕРЕКИСНОГО ОКИСЛЕНИЯ ЛИПИДА НА МИТОХОНДРИЮ ПЕЧЕНИ КРЫС ПРИ ПОСТНАТАЛЬНОМ ОНТОГЕНЕЗЕ //INTERNATIONAL JOURNAL OF DISCOURSE ON INNOVATION, INTEGRATION AND EDUCATION. – 2020. – Т. 1. – №. 5. – С. 78-86.
7. Niyazmetov B., Akhmedov R., Mirzaolimov M. UNCOU'LED RESPIRATION IN BIRD MITICHONDRIA: CONNECTION WITH THERMOGENESIS //Bulletin of Namangan State University: Vol. – 2019. – Т. 1. – №. 2. – С. 18.
8. Бохонова Н. С., Мирзаолимов М. М. ВЛИЯНИЕ КАЛОРИЙНО-ОГРАНИЧЕННОЙ ДИЕТЫ НА ПРОДОЛЖИТЕЛЬНОСТЬ ЖИЗНИ ЖИВОТНЫХ //ТА'ЛИМ VA RIVOJLANISH TANLILI ONLAYN ILMİY JURNALI. – 2022. – С. 139-142.
9. Анваров Ф. Р., Мирзаолимов М. М. ГЕРИАТРИЯ (ОПРЕДЕЛЕНИЕ ПОНЯТИЙ; ЗАДАЧИ, СТОЯЩИЕ ПЕРЕД ЭТИМ НАУКАМ; РАЗДЕЛЫ И ДОСТИЖЕНИЯ) //ТА'ЛИМ VA RIVOJLANISH TANLILI ONLAYN ILMİY JURNALI. – 2022. – С. 326-332.
10. Мирзаолимов М. М. и др. ГЕРИАТРИЯ (ОПРЕДЕЛЕНИЕ ПОНЯТИЙ; ЗАДАЧИ, СТОЯЩИЕ ПЕРЕД ЭТИМ НАУКАМ; РАЗДЕЛЫ И ДОСТИЖЕНИЯ) //ТА'ЛИМ VA RIVOJLANISH TANLILI ONLAYN ILMİY JURNALI. – 2022. – С. 143-149.
11. Таджибаева Г. И., Мирзаолимов М. М. КАЛАМУШЛАРДА СУРУНКАЛИ ЭМОЦИОНАЛ СТРЕСС МОДЕЛИНИ ЯРАТИШ ВА ОРГАНИЗМДАГИ БИОКИМЁВИЙ ЎЗГАРИШЛАРНИ АНИҚЛАШ //BARQARORLIK VA YETAKSHI TADQIQOTLAR ONLAYN ILMİY JURNALI. – 2022. – С. 166-170.
12. Mirzavalievich M. M., Adashaliyevich N. Q. Soy Protein, Isoflavones, and Cardiovascular Health //International Journal of Scientific Trends. – 2023. – Т. 2. – №. 4. – С. 10-18.
13. Атаханова С. Д. СТРЕСС ТАЪСИРИДА ОРГАНИЗМ АЪЗО ВА ТИЗИМЛАРИДАГИ ФИЗИОЛОГИК-БИОКИМЁВИЙ ЎЗГАРИШЛАР //BARQARORLIK VA YETAKSHI TADQIQOTLAR ONLAYN ILMİY JURNALI. – 2022. – Т. 2. – №. 4. – С. 215-219.
14. Сопиев Ш. К., Рўзибоева С. И., Мирзаолимов М. М. ЕДИНСТВО И ВЗАИМОСВЯЗЬ СТРУКТУРЫ СОРЕВНОВАТЕЛЬНОЙ ДЕЯТЕЛЬНОСТИ И СТРУКТУРЫ ПОДГОТОВЛЕННОСТИ //PEDAGOGS journali. – 2022. – Т. 10. – №. 4. – С. 220-228.
15. Таджибаева Г. И., Мирзаолимов М. М. МИТОХОНДРИЯНИНГ ТАРКИБИЙ ТУЗИЛИШИ ВА БАЖАРАДИГАН ВАЗИФАЛАРИ //PEDAGOGS journali. – 2022. – Т. 8. – №. 1.
16. Сопиев Ш. К., Рўзибоева С. И., Мирзаолимов М. М. ПОДХОДЫ К РАЗВИТИЮ ДЕТСКО-ЮНОШЕСКОГО СПОРТА И ПОСТРОЕНИЮ

МНОГОЛЕТНЕЙ ПОДГОТОВКИ И ПОСТРОЕНИЮ МНОГОЛЕТНЕЙ ПОДГОТОВКИ //PEDAGOGS journali. – 2022. – Т. 5. – №. 1. – С. 162-170.

17. Мирзаолимов М. М. и др. ГЕРИАТРИЯ (ОПРЕДЕЛЕНИЕ ПОНЯТИЙ; ЗАДАЧИ, СТОЯЩИЕ ПЕРЕД ЭТИМ НАУКАМ; РАЗДЕЛЫ И ДОСТИЖЕНИЯ) //TA'LIM VA RIVOJLANISH TANLILI ONLAYN ILMIY JURNALI. – 2022. – С. 143-149.

18. Таджибаева Г. И., Мирзаолимов М. М. КАЛАМУШЛАРДА СУРУНКАЛИ ЭМОЦИОНАЛ СТРЕСС МОДЕЛИНИ ЯРАТИШ ВА ОРГАНИЗМДАГИ БИОКИМЁВИЙ ЎЗГАРИШЛАРНИ АНИҚЛАШ //BARQARORLIK VA YETAKSHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 166-170.

19. Атаханова С. Д. СТРЕСС ТАЪСИРИДА ОРГАНИЗМ АЪЗО ВА ТИЗИМЛАРИДАГИ ФИЗИОЛОГИК-БИОКИМЁВИЙ ЎЗГАРИШЛАР //BARQARORLIK VA YETAKSHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – Т. 2. – №. 4. – С. 215-219.

20. Мирзаолимов М. М. и др. КИСЛОРОД ВА АЗОТ ФАОЛ ШАКЛЛАРИ СИНТЕЗИ //PEDAGOGS journali. – 2022. – Т. 7. – №. 1. – С. 394-400.

21. Рахматуллаев А. М., Мирзаолимов М. М. РОЛЬ НА-К-НАСОСА. АКТИВНЫЙ ТРАНСПОРТ ИОНОВ КАЛЬЦИЯ И ВОДОРОДА В КЛЕТКЕ //Journal of new century innovations. – 2023. – Т. 43. – №. 1. – С. 8-13.