

THE ROLE OF XENOBIANTS IN ENVIRONMENTAL DISTURBANCES AND THEIR IMPACT ON THE ORIGIN OF AUTOIMMUNE DISEASES.

KARIMOVA I.I., KUCHKAROVA L.S., TOKHIROVA SH.A., MAKHKAMOVA M., IBRAGIMOVA D.O’.

National University after named Mirzo Ulugbek.

In the second half of the 20th century, ecological problem is being considered to be one of the important problems. That uncomfortable condition begin effecting live organisms, namely human beings [2]. According to toxicity, lead among different xenobiotics is concluded to be sharp. After lead enters to organism of human, it begins affecting to diverse systems of organism by breaking down. For example, encephalopathy with lead appears in body when lead affects to cardiovascular system. In this situation, conductivity of capillaries increases [7]. 5-aminolevulin (ALA) inhibiting 5-aminolevulin dehydratase is produced, when amount of gem decreases. When radioactive ^{222}Rh is broke down, radionucleotides are produced. according to volume, izatops of lead account for 50% of radionucleotides and it leads to Saturnism in organism. Mainly, by breathing and enteral way lead from ecological objects is able to enter to the body. Toxicity from lead could form because of industry and condition of life [3; 7]. Following information was found out during some expirements; if large amount of lead is accumulated in body, arteriosklerotic changes could have tendency to develop in arteries of kindeys [5].

When if lead and kadmiy substance are influenced to rats, antioxidative activity of enzymes decrease, and it leads to oxidative stress form of lipids [9].

Saturnism diseases differ from together with variouse symptoms. Because of saturnism, variouse diseases appear in gastrointestinal tract and ability related to act also shows like these changes [6; 4].

Physiological circumstance of organism is able to influence absorbtion of salt with lead in body. Particularly, when human is hungry, absorbtion is responsible for 60% and when people are full up, this number is considersd to be 3%. However, large amount of lead is absorbed in proximal part of intestines and some lead is absorbed in duodenum [6].

Biotransformation of xenobiotics is accomplished by enzymes and these enzymes consist of 3 cycles: 1-cycle- sytochrome P450 is considered from monoxigenaza groupe; 2-cycle- glutation –S- transferase (GST) and arilomino –N-asetyltranferase; 3-cycle; glycoproteins take part in [1; 8].

As a result variouse xenobiotics influence body of human beings, several changes would develop in pancreas, by which diabetes is getting more and more important side. 1-type diabetes are considered to form by genetic changes, and in these, rats are not getting more and more better obesity and it is called NOD (non obese diabetic). And autoimmune diabetes are developed independently in these rats that were NOD [10; 11]

In expirements, rats are choosen as amodel and they isolated to 2 groupes. While physiologic solution was sent to 1-groupe, to second groupe alloxane was sent. Briefly, rats were isolated to positive and negative groups.

Changes in pancreatic α -amylasa in experimental diabetic animals and their correction with prenalone (M \pm m; n=6)

Groupes of rats.	Days of experiments.			
	Activiytiy of α -amylasa (gr/minute/gr.tissue.			
	5-day	10-day	20-day	30-day
Control	35,5 \pm 3,5	34,8 \pm 3,1	37,5 \pm 3,5	34,8 \pm 3,1
Experiment	53,2 \pm 3,6	52,3 \pm 2,9	51,2 \pm 3,3	46,1 \pm 2,6

According to expirements, activity of α -amylasa was determined homogenate of pancreas in every rat from all groups and α -amylasa's ratio was studied related to specific tissue .Activity of α -amylasa was increased respectively according to table above;49,9%, 50,3%, 36,5%.

In conclusion, α -amilasa' activity in diabetes with alloxane increases related to specific tissue. It is clear that the adaptation reaction is carried out by the hyperfunction of the synthesis of digestive enzymes under the experimental condition.

References:

1. Костюк С.А. Системабиотрансформации ксенобиотиков: гены детоксикации. Медицинские новости. №11.2020. С.12-16.
2. Желтиков А.А., Северин А.Е., Желтиков В.А., Кориеева Л.Н. Эколого-физиологический подход в стратегии снижения заболеваемости населения// Материалы X международного симпозиума «Эколого-физиологические проблемы адаптации». Москва, 2001.-С.188-189.
3. Здольник Т.Д. Токсиколого-гигиенические аспекты влияния металлов на функцию пищеварения// монография ; Ряз. гос. ун-т им. С.А. Есенина. - Рязань, 2007.- 172 с.
4. Ковальчук В.К., Иванова И.Л., Колдаев В.М. Роль окружающей среды в возникновении неинфекционных заболеваний пищеварительной системы в приморском крае// Гигиен.Санитар.журн.-Россия, 2011.-№3. -С.10-15.
5. Коробкова А.И., Сорокина Н.С., Молодкина Н.Н. Свинец и его действие на организм (обзор литературы)// Мед.труда. Россия, 2001. -№5.-С.29-34.
6. Лобанова Е.А., Сорокина Н.С., Семенова Л.С. Заболевания гастродуоденальной зоны у работающих в контакте со свинцом// Мед. труда.-Москва, 2001.-№ 5.-С.42-44.
7. Новикова М.А., Пушкарев Б.Г., Судаков Н.П., Никифоров С.Б., Гольдберг О.А. Явербаум П.М. Влияние хронической свинцовой интоксикации на организм человека // Сибирский медицинский журнал (Иркутск).2013-№2.-С.13-16.
8. Чурносов М.И. Чурносов, И.С. Полякова, С.П.Пахомов, В.С.Орлова. Молекулярные и генетические механизмы биотрансформации ксенобиотиков. 2011. С.
9. Pillai A., Gupta S. Antioxidant enzyme activity and lipid peroxidation in liver of female rats co –exposed to lead and cadmium: effects of vitamin E and Mn2+//Free radice research.-2005.-Vol.39(7).-P.707-712.
10. Andrea G., Carmen S., Silvina N., Rita R. Immunologic and genetic markers in insulin-dependentdiabetes mellitus in anargentine population. G. Andrea, S. Carmen, N. Silvina, R. Rita., 2001, P. 279-283.
11. Zhou Y., Liu S., Wang J., Yan X. Changes in blood glucose of elderly patients with gastric cancer combined with type 2 diabetes mellitus after radical operation and the effect of mediation adjustment for blood glucose on the recovery of gastric cancer // Oncol Lett. 2018. P. 4303-4308.