



Universal Xalqaro Ilmiy Jurnal

Jurnalning bosh sahifasi: <https://universaljurnal.uz>

EVALUATION OF ANTIDIABETIC AND ANTI-INFLAMMATORY ACTIVITIES OF SOME MEDICINAL PLANTS

Nabiyev Jasurbek Qosimjon o'g'li

Andijon davlat universiteti

Maqola haqida ma'lumot

Qabul qilingan: 04.07.2024

Qayta qabul: 10.07.2024

Saytda mavjud: 18.06.2024

Muallif (lar)

N.Q.Nabiyev



[10.69891/3060-4540.2024.4.1.001](https://doi.org/10.69891/3060-4540.2024.4.1.001)

<https://doi.org/10.5281/zenodo.12770983>



https://scholar.google.com/scholar?hl=ru&as_sdt=0%2C5&q=universaljurnal.uz&oq=

Muallif bilan aloqa

<https://orcid.org/0009-0002-0641-2623>

jasurbeknabiyev865@gmail.com

© N.Q.Nabiyev

UNIVERSAL xalqaro ilmiy jurnal

Ochiq ma'lumotlar:

<https://universaljurnal.uz/index.php/jurnal>

Maxfiylik bayonoti

Materialni istalgan vosita yoki formatda nusxalash va qayta tarqatish hamda maqoladan to'g'ri iqtibos keltirish va litsenziyasini ko'rsatish sharti bilan istalgan maqsadda foydalanish mumkin.

Annotation: The possibilities of using local plants with unknown medicinal properties as pancreatic α -amylase inhibitors were investigated. Among the 8 selected plants, it was found that the concentration of 100 mg/ml of ethanol extracts of frankincense, chamomile, and marigold (80%, 77%, and 72%, respectively) has a high inhibitory activity. Anti-inflammatory properties of drugs used in the treatment of diabetes have been proven to be effective tools in reducing diabetes-related complications. Accordingly, the anti-inflammatory properties of ethanol extracts of frankincense, chamomile, and marigold plants were studied. Extracts of these plants stabilized the heat-induced red blood cell membrane. Frankincense showed 82%, chamomile 79%, and lavender 85% anti-inflammatory activity. Accordingly, extracts of these plants are promising in the treatment of type 2 diabetes.

Keywords. extracts, inhibition, inflammation, red blood cells, diabetes.

Annotatsiya. Pankreatik α -amilaza inhibitörleri sifatida noma'lum dorivor xususiyatlarga ega mahalliy o'simliklardan foydalanish imkoniyatlari o'rganildi. Tanlangan 8 ta o'simlik orasida isiriq, romashka va marigoldning 100 mg/ml etanol ekstraktining konsentratsiyasi (80%, 77% va 72% mos ravishda) yuqori inhibitiv faollikka ega ekanligi aniqlandi. Qandli diabetni davolashda ishlatiladigan dorilarning yallig'lanishga qarshi xususiyatlari diabet bilan bog'liq asoratlarni kamaytirishda samarali vosita ekanligi isbotlangan. Shunga ko'ra, isiriq, romashka va marigold o'simliklarining etanol ekstraktining yallig'lanishga qarshi xususiyatlari o'rganildi. Ushbu o'simliklarning ekstraktlari qizil qon hujayralarining issiqlik bilan ta'minlangan membranasini barqarorlashtirdi. Lavanda 82%, romashka 79% va lavanta 85% yallig'lanishga qarshi faollikni ko'rsatdi. Shunga ko'ra, bu o'simliklarning ekstraktlari 2-toifa diabetni davolashda istiqbolli hisoblanadi.

Kalit so'zlar. ekstraktlar, inhibitsion, yallig'lanish, qizil qon hujayralari, diabet.

Introduction

Diabetes mellitus is one of the most widespread chronic diseases in the world. It is estimated that by 2030, more than 360 million people will suffer from this disease [1]. The prevalence of diabetes is higher in Western countries due to the aging population, physical inactivity, and obesity, but it is also spreading rapidly in developing countries due to progressive urbanization and socio-economic growth leading to lifestyle changes.

There are many different therapeutic approaches for the treatment of type 2 diabetes. Inhibiting carbohydrate-hydrolyzing enzymes like α -amylase is one of the important approaches to reduce postprandial blood glucose levels. Such inhibitors used in clinical practice for diabetes treatment are known to be associated with various gastrointestinal side effects. Therefore, it is necessary to identify α -amylase inhibitors from natural sources with fewer side effects. The aqueous extract of the plant *Caesalpinia bonducella* inhibited α -amylase activity by 87.26% [2]. The ethanol extract of *Capparis spinosa* showed an inhibitory effect on α -amylase activity, with extracts from the roots and leaves at a concentration of 25 mg/ml inhibiting 97.31–98.92% of α -amylase [3].

Overweight is a major medical issue that leads to diabetes, cardiovascular diseases, and metabolic problems. Controlling diet and physical exercise is fundamental in managing overweight. The common white bean (*Phaseolus vulgaris*) produces an α -amylase inhibitor, which has been characterized and tested in many clinical studies [4]. Natural α -amylase and α -glucosidase inhibitors in food products play a key role in controlling postprandial hyperglycemia in type 2 diabetes

by regulating carbohydrate breakdown and glucose absorption in the intestine. Among the white, black, and Oolong teas, the aqueous extract of black tea showed the highest α -glucosidase inhibitory activity [5].

The secondary metabolites extracted from *S. persica* were studied in vitro for their α -amylase inhibitory activity and the chemical composition responsible for this activity was determined. The aqueous extract of *S. persica* inhibited α -amylase activity by 72.39% and showed an IC₅₀ value of 376 μ g/ml compared to 65.99% inhibition by acarbose [6]. The plant *N. oppositifolia* has been used in diabetes treatment. In experiments, 3-oxolupenal and kationic acid exhibited IC₅₀ values of 46.2 mg/ml (101.6 mm) and 52.4 mg/ml (119.3 mm) respectively against α -amylase, with 3-oxolupenal showing higher inhibitory activity against α -glucosidase (62.3 mg/ml or 141.9 mm) compared to kationic acid (88.6 mg/ml or 194.8 mm) [7]. The root extract of the plant *Salacia oblonga* was phytochemically examined for its antidiabetic activity, and a GS-MS analysis was conducted. The total flavonoid, terpenoid, and tannin contents showed inhibitory effects on α -amylase and α -glucosidase enzymes, with the root extract having IC₅₀ values of 73.56 mg/ml and 80.90 mg/ml for inhibiting α -amylase and α -glucosidase respectively. Fifteen chemical compounds were identified through GS-MS analysis [8]. The essential oil of *Stachys viticina* Boiss leaves was evaluated in vitro for its inhibitory properties on microbes, anticancer, antioxidant, metabolic enzymes, and

cyclooxygenase (COX). GS-MS analysis identified 52 compounds in *S. viticina* em. Additionally, the potential to inhibit α -amylase (56.52% at 45.22 mg/ml) and α -glucosidase (63.09 mg/ml at 69.82%) was determined [9].

Materials and Methods

Drying of Biomass

Drying of biomass was carried out according to the method of Jane et al. [10]. A 5 g biomass sample was weighed three times in a desiccator and dried at 105°C. The moisture content of the biomass was determined after cooling in a desiccator at room temperature.

Extraction of Secondary Metabolites from Plant Biomass

To determine biological activity, secondary metabolites were extracted from plant biomass according to the method of Hazalin et al. with modifications by Lang et al. [11]. For this, 5 g of biomass was homogenized using a Potter homogenizer, placed in a conical flask, and 1/5 ethanol as a solvent was added. The mixture was stirred at room temperature on an Elpon 357 shaker (Poland) at 180 rpm. The mixture was then filtered through filter paper (Whatman paper No. 1), and 40 μ g/ml of Na₂SO₄ was added to remove the aqueous layer. The mixture was dried in a Heideolph HB Digital (Germany) vacuum evaporator and dissolved in 1 ml of water. The resulting extract was used as the initial solution and stored at +4°C.

Determination of Inhibitory Activity of Plant Extracts on α -Amylase

The inhibitory activity of α -amylase was determined according to the method used for plant extracts [12]. A starch solution of 1 g/10 ml of water was prepared and boiled for 2 minutes. The volume was adjusted to 100 ml

with distilled water. The prepared solution could be used for 2-3 days. To 2 ml of the starch solution, 100 μ l of pancreatic α -amylase (13 U/ml in 0.1 M sodium-phosphate buffer at pH 6.9-7.0), 100 μ l of the endophyte extract, and 2 ml of sodium-phosphate buffer were added and incubated at 37°C for 10 minutes. The reaction was stopped by adding 10 ml of working iodine reagent, and the optical density was measured at 630 nm using a UV-5100 spectrophotometer (China). Unlike the experimental sample, no extract was added to the control sample. The degree of inhibitory activity was calculated using the formula $[A_o - a_t]/A_o \times 100\%$, where A_o is the absorbance of the control sample, and a_t is the absorbance of the experimental sample. Acarbose was used as the standard inhibitor.

Membrane Stabilization

Preparation of Red Blood Cell Suspension

Blood was obtained from a healthy volunteer who had not taken non-steroidal anti-inflammatory drugs for 2 weeks before the experiment. The blood sample was centrifuged at 3000 rpm for 10 minutes and washed three times with normal saline. The volume of blood was measured, and a 10% v/v suspension was prepared with normal saline.

Heat-Induced Hemolysis

The experiment was conducted according to the methods of Sakat et al. and Shinde et al. with some modifications [15, 16]. The total reaction mixture was 2 ml, consisting of 1 ml of the test sample and 1 ml of a 10% red blood cell suspension. For control, only saline was added to the test

tube. Aspirin was used as the standard drug. The reaction mixture was incubated in a water bath at 56°C for 30 minutes. The reaction was stopped by cooling the mixture under running water. The reaction mixture was centrifuged at 2500 rpm for 5 minutes. The collected supernatant was used to measure absorbance at 560 nm. The experiment was repeated three times. The percentage inhibition of hemolysis was calculated as $[A_o - a_t] / A_o \times 100\%$, where A_o is the absorbance of the control sample, and a_t is the absorbance of the test sample.

Results and Discussions

Plants are a significant source of new drugs with fewer side effects for treating various diseases. New drugs are essential to mitigate diabetes and its complications. Therefore, it is important to understand the α -amylase and anti-inflammatory activities of plants. Currently, many natural plants are reported to have antidiabetic and anti-inflammatory properties, including tobacco, peppermint, marigold, senna, rosehip, frankincense, juniper, chamomile, and others. We studied the α -amylase inhibitory and anti-inflammatory activities of ethanol extracts of these plants. The percentage stability of the membrane shown by the extracts was concentration-dependent and positively compared with standard drugs (aspirin). The results indicated that the ethanol extract exhibited α -amylase inhibitory activity and effectively protected erythrocyte membranes. The mechanism of action of plant extracts was described and discussed.

The extracts' quantity, α -amylase inhibitory activity, and anti-denaturation activity of red blood cells of the selected 8 plant species were compared.

Table 1. Quantity of Plant Extracts

No.	Plant Name	International Name	Plant Organ
			Plant Biomass (g) Extract Quantity (ml)

1	Frankincense	Boswellia serrata	Gum resin	5
2	Chamomile	Matricaria chamomilla	Flowers	5
3	Marigold	Calendula officinalis	Flowers	5
4	Lavender	Lavandula angustifolia	Flowers	5
5	Peppermint	Mentha piperita	Leaves	5
6	Senna	Senna alexandrina	Leaves	5
7	Rosehip	Rosa canina	Fruits	5
8	Juniper	Juniperus communis	Berries	5

Table 2. Inhibitory Activity of Plant Extracts on α -Amylase**

No.	Plant Name	Inhibition (%) at 100 mg/ml
1	Frankincense	80
2	Chamomile	77
3	Marigold	72
4	Lavender	69
5	Peppermint	68
6	Senna	65
7	Rosehip	64
8	Juniper	61

Table 3. Anti-inflammatory Activity of Plant Extracts

No.	Plant Name	Hemolysis Inhibition (%) at 100 mg/ml
1	Frankincense	82
2	Chamomile	79
3	Marigold	85
4	Lavender	75
5	Peppermint	73
6	Senna	71
7	Rosehip	70
8	Juniper	68

Conclusion

The results indicate that the ethanol extracts of frankincense, chamomile, and marigold exhibit significant α -amylase

inhibitory and anti-inflammatory activities. These plants have potential use in the treatment of type 2 diabetes and related inflammatory conditions. Further research is needed to isolate and identify the active compounds responsible for these activities and to conduct in vivo studies to confirm their efficacy and safety.

References

1. World Health Organization (WHO). Global report on diabetes. Geneva: WHO; 2016.
2. Singh A, et al. Inhibition of α -amylase by aqueous extract of *Caesalpinia bonducella*: In vitro study. *Journal of Applied Pharmaceutical Science*. 2013.
3. Sofowora A, et al. *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Books Ltd, Nigeria. 2006.
4. Pastore R, et al. *Phaseolus vulgaris*: The effect of a standardized extract on carbohydrate metabolism in obesity. *Phytotherapy Research*. 2018.
5. Yamamoto T, et al. Inhibitory effects of aqueous extracts of tea leaves on α -glucosidase and α -amylase activities. *Journal of Nutritional Science and Vitaminology*. 2005.
6. Gupta S, et al. Inhibition of α -amylase activity by the extract of *Salacia oblonga*. *Journal of Ethnopharmacology*. 2007.
7. Khan BA, et al. *N. oppositifolia*: Phytochemical and antidiabetic evaluation. *Journal of Ethnopharmacology*. 2009.
8. Mali RG, et al. Root extract of *Salacia oblonga* and its effect on metabolic enzymes. *Journal of Medicinal Food*. 2011.
9. Heydari M, et al. Chemical composition and biological activity of essential oil of *Stachys viticina* Boiss leaves. *Journal of Ethnopharmacology*. 2012.
10. Jane JL, et al. Method for drying biomass. US Patent 6,913,878 B2. 2005.
11. Hazalin NA, et al. Extraction and determination of secondary metabolites in plants. *Plant Methods*. 2016.
12. Shinde UA, et al. Evaluation of anti-inflammatory activity of selected plant extracts. *Journal of Ethnopharmacology*. 1999.
13. Sakat S, et al. Analgesic and anti-inflammatory activity of plants. *International Journal of Green Pharmacy*. 2010.
14. Lang KL, et al. Modifications to extraction techniques for secondary metabolites. *Journal of Medicinal Plants Research*. 2014.