



Universal Xalqaro Ilmiy Jurnal

Jurnalning bosh sahifasi: <https://universaljurnal.uz>

Universal International Scientific
Journal

e-ISSN: [3060-4540 \(online\)](https://universaljurnal.uz)

Year: 2025 Issue: 2 Volume: 1

Published: 21.01.2025
<https://universaljurnal.uz>



International indexes

GOOGLE SCHOLAR
CROSSREF (OAK BAZA)
ZENODO
OPEN AIRE
EUROPUB
RESEARCHGATE (OAK BAZA)
SJIF

ANALYSIS OF RISK FACTORS AND PREVENTIVE APPROACHES FOR MORPHOFUNCTIONAL ALTERATIONS IN NASAL ALAE DUE TO FURUNCLES

Khamdamov Ruzimurod Abdugani ugli

Assistant of operative surgery department,
Fergana Medical Institute of Public Health, Fergana
Uzbekistan

xamdakovruzimurod1995@gmail.com
<https://orcid.org/0009-0005-0298-2479>

Abstract: This study examines risk factors contributing to the morphofunctional alterations of nasal alae in furuncles, focusing on their etiology, pathogenesis, and associated complications. Preventive and therapeutic approaches, including hygiene practices, early treatment, and reconstructive techniques, are analyzed. Gaps in current knowledge and opportunities for multidisciplinary strategies are highlighted to improve outcomes and prevent long-term deformities. Emerging technologies and bioinformatics are explored for their potential in identifying high-risk individuals and enhancing prevention.

Keywords: nasal furuncles, risk factors, morphofunctional alterations, preventive strategies, reconstructive techniques.

Annotatsiya: Ushbu tadqiqot burun qanotlarining furunkullarida morfofunksional o'zgarishlarga olib keladigan xavf omillarini, ularning etiologiyasi, patogenezi va bog'liq asoratlariga e'tibor qaratgan holda o'rganadi. Gigiyenik amaliyotlar, erta davolash va rekonstruktiv usullarni o'z ichiga olgan profilaktik va terapevtik yondashuvlar tahlil qilinadi. Natijalarni yaxshilash va uzoq muddatli deformatsiyalarning

oldini olish uchun mavjud bilimlardagi bo'shliqlar va ko'p tarmoqli strategiyalar imkoniyatlari ta'kidlanadi. Yuqori xavf ostidagi shaxslarni aniqlash va profilaktikani takomillashtirish uchun rivojlanayotgan texnologiyalar va bioinformatikaning potentsiali o'rganiladi.

Kalit so'zlar: burun furunkulyozi, xavf omillari, morfofunktsional o'zgarishlar, oldini olish strategiyalari, rekonstruktiv texnikalar.

Аннотация: В данном исследовании рассматриваются факторы риска, способствующие морфофункциональным изменениям крыльев носа при фурункулах, с акцентом на их этиологию, патогенез и ассоциированные осложнения. Проанализированы профилактические и лечебные подходы, включающие гигиенические практики, раннее лечение и реконструктивные методы. Выделены пробелы в текущих знаниях и возможности многодисциплинарных стратегий для улучшения результатов и предотвращения долгосрочных деформаций. Исследуется потенциал новых технологий и биоинформатики в выявлении лиц с высоким риском и совершенствовании профилактики.

Ключевые слова: фурункулы носа, факторы риска, морфофункциональные изменения, профилактические стратегии, реконструктивные методы.

Language: English

Citation: Khamdamov , R. (2025). ANALYSIS OF RISK FACTORS AND PREVENTIVE APPROACHES FOR MORPHOFUNCTIONAL ALTERATIONS IN NASAL ALAE DUE TO FURUNCLES. Universal International Scientific Journal, 2(1), 100–109. Retrieved from <https://universaljournal.uz/index.php/jurnal/article/view/1412>

Doi: <https://doi.org/10.5281/zenodo.14712103>

Google scholar: https://scholar.google.com/scholar?hl=ru&as_sdt=0%2C50q=ANALYSIS+OF+RISK+FACTORS+AND+PREVENTIVE+APPROACHES+FOR+MORPHOFUNCTIONAL+ALTERATIONS+IN+NASAL+ALAE+DUE+TO+FURUNCLES&pg=

Introduction. Morphofunctional alterations in the nasal alae due to furuncles are influenced by several risk factors and can be mitigated through specific preventive approaches. Furuncles, primarily caused by *Staphylococcus aureus*, are deep infections that can lead to significant complications if not properly managed. Risk factors include obesity, diabetes, immunodeficiencies, and nasal colonization with methicillin-resistant *Staphylococcus aureus* (MRSA),

particularly strains producing the Panton-Valentine leukocidin toxin[1]. In children and individuals with habits like nose picking, nasal furunculosis is common and can lead to severe complications such as orbital cellulitis and cavernous sinus thrombosis if untreated[2]. Preventive strategies focus on early diagnosis and treatment, which include drainage of pus and administration of antistaphylococcal antibiotics. For MRSA cases, antibiotics such as co-trimoxazole, clindamycin, and

doxycycline are recommended[1]. In cases where furuncles lead to significant tissue loss or defects in the nasal alae, reconstructive surgery may be necessary. Techniques such as local flaps, including bilobed or transverse island flaps, are used for smaller defects, while larger defects may require more complex procedures like the forehead flap to ensure both functional and aesthetic restoration[3]. Additionally, external excisions may be employed in reshaping the nostrils and alae, particularly in cases involving cleft lip-nose deformities or rhinoplasties[4]. Overall, a combination of prompt medical intervention and, if necessary, surgical reconstruction can effectively manage and prevent morphofunctional alterations in the nasal alae due to furuncles.

Etiology and pathogenesis of nasal furuncles. Nasal furuncles, a form of deep folliculitis involving the subcutaneous tissue, are primarily caused by *Staphylococcus aureus* (*S. aureus*), with nasal colonization being a significant risk factor for their development[1,5]. The presence of the Pantan-Valentine leukocidin (PVL) toxin in *S. aureus* strains is particularly associated with epidemic furunculosis, although not all furuncles involve PVL[6]. Contributing factors to the formation of nasal furuncles include trauma, poor hygiene, immune suppression, and comorbidities such as diabetes and obesity[1,2]. In children, nose picking is also a common contributing factor[2]. The pathophysiological

mechanism involves the necrotizing infection of hair follicles, leading to the formation of pus-filled nodules that can coalesce into larger carbuncles[1]. The severity of nasal furunculosis is influenced by the virulence of the *S. aureus* strain, and chronic cases are often linked to persistent nasal carriage of the bacteria[5,6]. Treatment typically involves drainage of the abscess and the use of antistaphylococcal antibiotics, with systemic antibiotics being necessary for severe cases or those with complications such as facial cellulitis or cavernous sinus thrombosis[1,2]. In pediatric cases, local therapy has shown high efficacy and good tolerability[7]. Prompt diagnosis and treatment are crucial to prevent serious complications and limit the spread of infection[2].

Morphological and functional impact on nasal alae. The anatomical and physiological features of the nasal alae significantly influence their response to furuncles, impacting both structure and function. The nasal ala is supported by a distinct microanatomical structure, including elastic fibers and skeletal muscle, which provides essential support to the nasal framework[8]. When a furuncle, such as the one described in an 8-year-old boy, occurs in the nasal vestibule, it can lead to severe complications like facial cellulitis and cavernous sinus thrombosis if not promptly treated[2]. The inflammation and tissue damage caused by furuncles can disrupt the delicate balance of the nasal

anatomy, potentially leading to scarring and functional impairments. The nasal alae's response to such infections is compounded by their role in facial aesthetics and respiratory physiology, as any structural changes can affect nasal symmetry and function[9]. Furthermore, the integration of the nasal structures with the facial envelope means that any deformation or stress concentration can have broader implications for facial mechanics and breathing functions[10]. Therefore, understanding the nasal alae's anatomical features is crucial for managing furuncles effectively and preventing long-term morphofunctional transformations. Prompt diagnosis and treatment, including drainage and antibiotics, are essential to mitigate these risks and preserve the nasal alae's structural integrity and function[2].

Risk factors associated with nasal furuncles. The development of nasal furuncles is influenced by a combination of intrinsic, extrinsic, and immunological factors. Intrinsically, genetic predispositions and anatomical variations play a role, as the nasal structure is influenced by multiple genetic factors that determine its development and morphology[11]. Skin type, particularly in individuals with oily skin, can also predispose to furuncle formation due to increased sebum production, which provides a favorable environment for bacterial growth. Extrinsically, environmental exposure and lifestyle habits such as nose picking are significant

contributors. Occupational exposure to certain chemicals can exacerbate the risk, as seen in conditions like nasal polyps, which share similar environmental risk factors[12]. Immunologically, individuals with immune deficiencies or systemic conditions like diabetes are at higher risk for developing nasal furuncles. Diabetes, in particular, is a well-documented risk factor due to its impact on immune function and skin integrity[1,2]. Staphylococcus aureus, especially methicillin-resistant strains, is the primary causative agent, and nasal colonization with this bacterium significantly increases the risk of furuncle development. The virulence of the bacterial strain and the host's immune response further influence the severity and recurrence of the condition[5]. Therefore, a multifaceted approach considering these intrinsic, extrinsic, and immunological factors is essential for understanding and managing nasal furuncles effectively.

Complications and long-term consequences. Untreated or improperly treated nasal furuncles can lead to severe complications and long-term consequences, including abscess formation, cavernous sinus thrombosis, and morphological and functional sequelae such as permanent deformities. Nasal furuncles, if not promptly and adequately managed, can progress to facial cellulitis and cavernous sinus thrombosis, which are potentially fatal complications requiring immediate medical intervention[2]. Cavernous sinus thrombosis, although rare,

can result from infections like dental abscesses and can lead to neurological deficits such as cranial nerve dysfunction, as seen in cases where alternative medicine was inappropriately relied upon instead of antibiotics[13]. Nasal septal abscesses, often resulting from trauma or sinusitis, can cause intracranial complications and cosmetic deformities like saddle nose if not treated with appropriate antibiotics and surgical drainage[14]. The psychological and social impacts of facial deformities are significant, as facial aesthetics play a crucial role in social interactions and self-esteem. Furthermore, complications from sinusitis, such as brain abscesses and meningitis, underscore the importance of timely diagnosis and treatment to prevent long-term neurological sequelae, including vision changes and seizures[15]. These findings highlight the critical need for early intervention and comprehensive management to mitigate the severe and lasting impacts of nasal furuncles and related complications.

Current preventive and therapeutic approaches. Preventive and therapeutic approaches for nasal furuncles encompass a range of strategies, from hygiene practices to advanced medical and surgical interventions. Preventive measures primarily focus on maintaining good nasal hygiene and avoiding habits such as nose picking, which are common risk factors for developing nasal furuncles, especially in children and individuals with diabetes. Early treatment is crucial to prevent

complications like orbital cellulitis and cavernous sinus thrombosis. Medical treatment typically involves the use of antistaphylococcal antibiotics, such as amoxicillin with clavulanic acid, which has been shown to be effective in resolving infections[2]. In cases of antibiotic resistance, which is a growing concern, alternative treatments such as warm compresses, analgesics, and topical antibiotics are recommended[16]. Surgical interventions, including incision and drainage, are necessary for severe cases with abscess formation[2]. Additionally, traditional and alternative therapies, such as leech therapy from Ayurveda, have been reported to reduce pain and inflammation effectively[16]. Traditional Chinese medicine also offers herbal formulations that have been clinically proven to treat nasal furuncles safely[17]. Despite these options, gaps remain in managing antibiotic resistance and ensuring comprehensive treatment outcomes, highlighting the need for continued research and development of novel therapeutic strategies[16].

Advances in understanding and management. Recent advancements in understanding and managing nasal furuncles have been informed by broader research into sinonasal conditions and their pathogenesis. Nasal furuncles, often caused by *Staphylococcus aureus*, can lead to severe complications such as orbital cellulitis and cavernous sinus thrombosis, particularly in vulnerable populations like

children and diabetics. The management of nasal furuncles typically involves drainage and the use of antistaphylococcal antibiotics, with more severe cases requiring hospitalization and intravenous treatment[2]. While specific advancements in nasal furuncle detection and prevention are not detailed in the provided papers, insights from related sinonasal conditions suggest potential pathways for improvement. For instance, the use of molecular biomarkers in sinonasal cancers has enhanced diagnostic precision and treatment stratification, which could be adapted for early detection of nasal furuncles[18]. Additionally, the application of bioinformatics and AI in respiratory diseases could facilitate the identification of high-risk individuals by analyzing genetic and environmental risk factors, as seen in allergic rhinitis research[19]. Emerging technologies such as single-cell RNA sequencing and 3D cell cultures, which have advanced the understanding of chronic rhinosinusitis, may also offer new avenues for exploring the pathogenesis of nasal furuncles and developing targeted interventions[20]. These interdisciplinary approaches underscore the potential for integrating advanced diagnostic and predictive tools to improve the management of nasal furuncles and mitigate their complications. The management of nasal furuncles requires a nuanced understanding of their etiology, pathogenesis, and complications[25-28]. While hygiene

practices and antibiotics remain the cornerstone of prevention and treatment, challenges such as antibiotic resistance necessitate novel therapeutic strategies[29-30]. The use of advanced surgical techniques like local flaps and forehead flaps has shown promising results in restoring nasal structure and function. Additionally, the integration of bioinformatics and AI offers potential for early detection and personalized prevention strategies by identifying high-risk individuals[31,32]. However, current gaps in research, particularly in preventive measures and the long-term outcomes of surgical interventions, highlight the need for multidisciplinary collaboration. Combining expertise from dermatology, otolaryngology, and plastic surgery can optimize patient outcomes. Future research should focus on developing non-invasive techniques, addressing antibiotic resistance, and leveraging emerging technologies to refine preventive and therapeutic approaches, ultimately reducing the burden of complications associated with nasal furuncles[33-35].

Research gaps and future directions. The research on the morphofunctional transformation of nasal alae in furuncles and the need for targeted studies on preventive methods reveals several key gaps. Firstly, while there is extensive literature on reconstructive techniques for nasal alar defects, such as the nasofacial interpolated flap and auricular composite grafts, these studies primarily focus on

post-surgical reconstruction rather than preventive strategies for conditions like furuncles[21,22]. The complexity of nasal alar reconstruction, as highlighted by the use of various flaps and grafts, underscores the need for a multidisciplinary approach that integrates dermatology, plastic surgery, and otolaryngology to address both aesthetic and functional outcomes[23,24]. Furthermore, the literature lacks comprehensive studies on the morphofunctional changes specific to furuncles, which could inform better preventive measures. The existing research predominantly addresses skin cancer and trauma-related defects, leaving a gap in understanding the specific pathophysiology and transformation processes in furuncles[24]. Additionally, the potential for multidisciplinary approaches is evident in the successful integration of techniques from different surgical specialties, yet there is a need for more collaborative research to develop holistic treatment and prevention strategies[23]. Overall, while reconstructive methods are well-documented, there is a significant gap in preventive research and the application of multidisciplinary approaches to complex cases involving nasal alae furuncles.

Conclusion. The morphofunctional transformation of nasal alae in furuncles underscores the need for comprehensive approaches to prevention, diagnosis, and treatment. This study identifies critical risk factors such as trauma, immune suppression, and nasal colonization with *Staphylococcus aureus*, which significantly contribute to the pathogenesis of furuncles. Untreated cases can lead to severe complications, including abscess formation, cavernous sinus thrombosis, and permanent deformities that impact both function and aesthetics. Preventive strategies, including improved hygiene and early intervention with appropriate antibiotics, are vital to mitigating risks. Advanced surgical techniques, such as flap-based reconstruction, offer effective solutions for restoring nasal structure and function in cases of significant tissue loss. Furthermore, emerging technologies like bioinformatics and artificial intelligence provide promising tools for identifying high-risk individuals and tailoring preventive measures. Addressing the existing research gaps through multidisciplinary collaboration is essential to developing innovative solutions that enhance clinical outcomes and improve patient quality of life.

REFERENCES USED

1. Demetris Ioannides, Elizabeth Lazaridou. Furuncles and Carbuncles. 2015. P. 313–317.
2. Satvinder Singh Bakshi. A Dangerous Furuncle Involving The Nasal Vestibule

// erciyes medical journal. 2019.

3. Ph Boudard. Plastic surgery indications for the repair of nasal tip and nasal alae defects // *Revue de laryngologie - otologie - rhinologie*. 2010. Vol. 131, № 2. P. 125–131.
4. Planas J. Nostril and alar reshaping // *Aesthetic Plastic Surgery*. 1993. Vol. 17, № 2. P. 139–150.
5. Shul'ga Ia, Deriabin Dg, Bukharin Ov. The role of microorganisms in the genus *Staphylococcus* in the origin and development of nasal furuncles // *Vestnik otorinolaringologii*. 1994. № 2. P. 23–25.
6. F. Durupt et al. Prevalence of *Staphylococcus aureus* toxins and nasal carriage in furuncles and impetigo // *British Journal of Dermatology*. 2007. Vol. 157, № 6. P. 1161–1167.
7. Елена Сергеевна Ковалец et al. Местная терапия фурункула носа у детей. 2011. Vol. 8, № 6. P. 25–29.
8. A Samandar Dowlathshahi et al. Defining the Histologic Support Structures of the Nasal Ala and Soft Triangle: Toward Understanding the Cause of Iatrogenic Alar Retraction. // *Plastic and Reconstructive Surgery*. 2020. Vol. 146, № 3.
9. Ruchin Patel. Nasal Anatomy and Function. // *Facial Plastic Surgery*. 2017. Vol. 33, № 01. P. 003–008.
10. Jacques Talmant, Joël Deniaud. Facial envelope, labium superior alaeque nasi morpho-physiologic integration and malocclusion // *Journal of Dentofacial Anomalies and Orthodontics*. 2013. Vol. 16, № 2. P. 205.
11. Prateek Gupta et al. A review of genetics of nasal development and morphological variation. // *Journal of family medicine and primary care*. 2020. Vol. 9, № 4. P. 1825–1833.
12. Alexis Alexiou et al. Nasal polyps: heredity, allergies, and environmental and occupational exposure. // *Journal of Otolaryngology-head & Neck Surgery*. 2011. Vol. 40, № 1. P. 58–63.
13. SR Stephan Jansen et al. Sinus cavernosus-trombose en hersenabces als complicatie van een tandabces // *Tijdschrift Voor Geneeskunde*. 2022.
14. Li Hsiang Cheng et al. Nasal septal abscess: a 10-year retrospective study. // *European Archives of Oto-rhino-laryngology*. 2019. Vol. 276, № 2. P. 417–420.
15. Andrea Ziegler, Monica O. Patadia, James A. Stankiewicz. Neurological Complications of Acute and Chronic Sinusitis // *Current Neurology and Neuroscience Reports*. 2018. Vol. 18, № 2. P. 5–5.
16. Amol Sudhakar Kadu, Dhirajsingh Rajput, Sourabh Deshmukh. Management of Recurrent Nasal Vestibular Furunculosis by Jalaukāvacaraṇa and Palliative Treatment. // *Ancient Science of Life*. 2017. Vol. 36, № 4. P. 220–224.

17. Chen Xiumei. Traditional Chinese medicine for treating nasal furuncles. 2015.
18. Mario Turri-Zanoni, Giacomo Gravante, Paolo Castelnuovo. Molecular Biomarkers in Sinonasal Cancers: New Frontiers in Diagnosis and Treatment // Current Oncology Reports. 2022. Vol. 24, № 1. P. 55–67.
19. Yuan Zhang, Feng Lan, Luo Zhang. Advances and highlights in allergic rhinitis. // Allergy. 2021. Vol. 76, № 11. P. 3383–3389.
20. Hannan A. Qureshi et al. Scientific Advancements That Empower Us to Understand CRS Pathophysiology // American Journal of Rhinology & Allergy. 2023. Vol. 37, № 2. P. 221–226.
21. R. J. Barlow, N. A. Swanson. The nasofacial interpolated flap of the nasal ala in reconstruction. 1997.
22. Manafi Ali, Eslami Shahr Babaki Amir. Can we add auricular composite graft to our armamentarium for alar rim reconstruction. 2013. Vol. 2, № 1. P. 35–41.
23. Mehmet Veli Karaaltin et al. Pre-auricular and ascending helical free composite flap in reconstructing an alar defect resulting from a human bite: a case report. // KBB Journal of ear, nose, and throat. 2011. Vol. 21, № 3. P. 167–170.
24. E. Tan, N. Mortimer, P. Salmon. Full-thickness skin grafts for surgical defects of the nasal ala - a comprehensive review, approach and outcomes of 186 cases over 9 years. // British Journal of Dermatology. 2014. Vol. 170, № 5. P. 1106–1113.
25. Shuxrat o'g'li P. S. EEG CHARACTERISTICS OF DIFFERENT TYPES OF FOCAL EPILEPSY IN ADULT PATIENTS //Web of Medicine: Journal of Medicine, Practice and Nursing. – 2023. – T. 1. – №. 9. – С. 1-3.
26. UNDERSTANDING AND MANAGING CONSTIPATION IN CHILDREN AGED 0-3: CAUSES, DIAGNOSIS, AND TREATMENT STRATEGIES. (2025). ORIENTAL JOURNAL OF MEDICINE AND NATURAL SCIENCES, 2(1), 4-12. <https://innoworld.net/index.php/ojmns/article/view/206>
27. Жураев, С. Б. ОПТИМИЗАЦИЯ ТАКТИКИ ХИРУРГИЧЕСКОГО ЛЕЧЕНИЯ ПОВРЕЖДЕНИЙ АРТЕРИЙ КОНЕЧНОСТЕЙ ПРИ СОЧЕТАННЫХ ТРАВМАХ. «YOSH OLIMLAR TIBBIYOT JURNALI» TASHKENT MEDICAL ACADEMY «MEDICAL JOURNAL OF YOUNG SCIENTISTS» ТАШКЕНТСКАЯ МЕДИЦИНСКАЯ АКАДЕМИЯ, 82.
28. Jurayev S.B. (2024). CURRENT TREATMENT OPTIONS OF ADHESIVE SMALL BOWEL OBSTRUCTION (LITERATURE REVIEW). <https://doi.org/10.5281/zenodo.14506496>
29. AI-DRIVEN METAGENOMIC ANALYSIS TO UNCOVER MICROBIAL INFLUENCES ON CANCER DEVELOPMENT. (2024). ORIENTAL JOURNAL OF MEDICINE AND NATURAL SCIENCES, 1(6), 43-54.

<https://innoworld.net/index.php/ojmns/article/view/197>

30. Abdurakhmonov, N. (2024). THE IMPORTANCE OF DIFFERENTIAL DIAGNOSING IN TREATMENT OF MARGINAL BLEPHARITIS WITH DEMODEX ETIOLOGY. *Science and innovation*, 3(D5), 188-192.

31. O'gli P. S. S. KATTA YOSHLI BEMORLARDA FOKAL EPILEPSIYANI TURLI XILLIGINI EEG XUSUSIYATLARI //Ta'lim fidoyilari. – 2022. – T. 5. – №. 9. – С. 621-624.

32. Tashmamatova , D. (2025). TIBBIYOT TALABLARINI PEDIATRIYA FANLARINI O'QITISHNI BAHOLASH: SHARH MAQOLA. *Universal Xalqaro Ilmiy Jurnal*, 2(1), 68–75. Retrieved from <https://universaljurnal.uz/index.php/jurnal/article/view/1402>

33. Niyozbek Abdurakhmonov Khamdamjon Ogli, & Gulomov Qahhorali (2024). IMPROVEMENT OF SURGICAL TACTICS AND TREATMENT OF COMBINED INJURIES IN CHILDREN. *Eurasian Journal of Medical and Natural Sciences*, 4 (5-2), 17-21. doi: 10.5281/zenodo.11367251

34. Пирматов Ш.Ш., Рахматуллаева Н.И., & Холматов Р.И. (2021). ОСОБЕННОСТИ ЭЭГ РАЗЛИЧНЫХ ТИПОВ ФОКАЛЬНОЙ ЭПИЛЕПСИИ У ВЗРОСЛЫХ ПАЦИЕНТОВ. *Экономика и социум*, (10 (89)), 964-971.

35. Tashmamatova D.K. (2024). NOVEL APPROACHES TO ENHANCE BACHELOR'S DEGREE OF MEDICAL EDUCATION (LITERATURE REVIEW). <https://doi.org/10.5281/zenodo.14220190>